

Local existence and uniqueness for a semi-linear accretive wave equation

H. FAOUR*, A. Z. FINO¹ AND M. JAZAR¹

June 18, 2010

Abstract

We study local existence and uniqueness in the phase space $H^\mu \times H^{\mu-1}(\mathbb{R}^N)$ of the solution of the semilinear wave equation $u_{tt} - \Delta u = u_t |u_t|^{p-1}$ for $p > 1$.

Keywords: Wave equations; Strichartz estimates.

1 Introduction and main results

A very rich literature has been done on the semi-linear wave equation

$$u_{tt} - \Delta u = au_t |u_t|^{p-1} + bu |u|^{q-1}$$

with a, b, p and q are real numbers, $p, q \geq 1$. When $a \leq 0$ and $b = 0$ then the damping term $au_t |u_t|^{p-1}$ ensures global existence in time for arbitrary data (see, for instance, Harraux and Zuazua [8] and Kopackova [10]). When $a \leq 0, b > 0$ and $p > q$ or when $a \leq 0, b > 0$ and $p = 1$ then one can cite, for instance, Georgiev and Todorova [5] and Messaoudi [12], that show the existence of global solutions under negative energy condition.

The first to consider the case $a > 0$ was Haraux [7] (with $b = 0$ on bounded domain), who construct blowing up solutions for arbitrary small initial data. See also Jazar and Kiwan [?] and the references therein for the same equation on bounded domain.

*Cermics, Paris-Est-ENPC, ParisTech, 6 et 8 avenue Blaise Pascal, Cité Descartes Champs-sur-Marne, 77455 Marne-la-Vallée Cedex 2, France. E-mail: faourh@cermics.enpc.fr.

¹A. Z. Fino and M. Jazar, LaMA-Liban, AZM Research Center, Lebanese University, P.O. Box 37, Tripoli Lebanon. E-mails: afino@ul.edu.lb, mjazar@ul.edu.lb.

The first is supported by the project ANR MICA (2006-2010).

The second author is supported by a grant from the Lebanese National Research Council.

The third authors is supported by a grant from the Lebanese University.

In this paper we consider the case $a = 1$ and $b = 0$, i.e. the semi-linear accretive wave equation

$$\begin{cases} u_{tt} - \Delta u = u_t |u_t|^{p-1} & x \in \mathbb{R}^N, t > 0, \\ u(0, x) = u_0(x), \quad u_t(0, x) = u_1(x) & x \in \mathbb{R}^N. \end{cases} \quad (1)$$

To our knowledge, no local existence result was done for this equation. The phase space to consider here is $Y^\mu := H^\mu \times H^{\mu-1}(\mathbb{R}^N)$, and we are looking to find conditions on the nonlinearity p and the order μ so that we have local existence. Due to our method, based on the use of Strichartz estimates (Proposition 3) and bounds on a power of a function in a Sobolev space $\|h^p\|_{H^s}$ by the norm of the initial function $\|h\|_{H^r}$, we need that p or μ to be integer. This is done in the following two theorems

Theorem 1 (*Case p integer*)

Let $p \in \mathbb{N} \setminus \{0, 1\}$ and $\mu \in \mathbb{R} \cap [1, \infty)$ such that

$$\begin{cases} 1 < p < \infty & \text{if } \mu \geq 1 + \frac{N}{2}, \\ 1 < p < \frac{N+4-2\mu}{N+2-2\mu} & \text{if } 1 \leq \mu < 1 + \frac{N}{2}. \end{cases} \quad (2)$$

For $(u_0, v_0) \in Y^\mu$, there exists a maximal time $T_{\max} > 0$ and a unique solution $(u, u_t) \in C^0([0, T_{\max}); Y^\mu)$ of problem (1). Moreover, if $T_{\max} < \infty$, we have $\|u(t)\|_{H^\mu} + \|u_t(t)\|_{H^{\mu-1}} \rightarrow \infty$ as $t \rightarrow T_{\max}$.

One can compare to the case $a = 0$ and $b = 1$, the restriction on p is the same by taking $\mu = 2$, see for instance [11, 14]).

In the previous theorem, p is integer. In the following theorem this is no longer the case. However, the dimension must be less than 3 or equal. This is due to Proposition 4 in which, we obtain L^∞ -estimates on the wave kernel. In what follows denote by $Y^{2,\infty} := W^{2,\infty} \times W^{1,\infty}(\mathbb{R}^N)$.

Theorem 2 (*Case p real*)

Let $1 \leq N \leq 3$, $\mu \in (1, 2) \cap \mathbb{N}^*$ and $p \in (1, \infty) \cap [\mu - 1, \infty)$. Then for all $(u_0, v_0) \in Y^\mu \cap Y^{2,\infty}$ there exists $T > 0$ and a unique solution $(u, u_t) \in C^0([0, T]; (L^\infty \cap H^\mu) \times L^\infty \cap H^{\mu-1}(\mathbb{R}^N))$ of (1).

2 Preliminary notations and results

In this section we use the notations used by [6]. For $x \in X$ a normed vector space we denote by $\|x; X\|$ the norm of x , and for $(x, y) \in X \times Y$ then, naturally, $\|(x, y); X \times Y\| = \|x; X\| + \|y; Y\|$. Finally, for $q \in [1, +\infty)$

define the norm $\|f; L^q(0, T; X)\|^q := \int_0^T \|f(t); X\|^q dt$ with the usual one for $q = +\infty$.

Consider the inhomogeneous wave equation in $\mathbb{R} \times \mathbb{R}^N$

$$\begin{cases} u_{tt} - \Delta u = f \\ u(0, x) = u_0(x), \quad u_t(0, x) = v_0(x). \end{cases} \quad (3)$$

We define the operator $\sigma := (-\Delta)^{1/2}$, which could be defined as $\sigma u(x) = \mathcal{F}^{-1}(|\xi| \mathcal{F}(u)(\xi))(x)$, and $K(t) := \sigma^{-1} \sin \sigma t$, $\dot{K}(t) := \cos \sigma t$. The solution of (3) could be written as $u = \theta + \omega$, where θ is the solution of the homogeneous equation with the same initial data

$$\begin{cases} \theta_{tt} - \Delta \theta = 0 \\ \theta(0, x) = u_0(x), \quad \theta_t(0, x) = v_0(x) \end{cases} \quad (4)$$

namely

$$\theta(t) = \dot{K}(t)u_0 + K(t)v_0 \text{ and } \theta_t(t) = \Delta K(t)u_0 + \dot{K}(t)v_0$$

that we denote by $H(t)U_0 = (\theta(t), \theta_t(t))$ where $U_0 := (u_0, v_0)$. And ω is the solution of the inhomogeneous equation with zero initial data

$$\begin{cases} \omega_{tt} - \Delta \omega = f \\ \omega(0, x) = \omega_t(0, x) = 0. \end{cases} \quad (5)$$

The solution of (5) could be written, for $t \geq 0$, as

$$\omega(t) = \int_0^t K(t-s)f(s)ds = K \star f(t) \text{ and } \omega_t(t) = \int_0^t \dot{K}(t-s)f(s)ds = \dot{K} \star f(t).$$

The initial data U_0 will be taken in the phase space Y^μ for $\mu \in \mathbb{R}$ where H^μ is the homogeneous Sobolev space (See [16]). We will use the following “simplified version” of the generalized Strichartz inequality [15]:

Proposition 3 (*Proposition 3.1 of [6]*) *Let $\rho_1, \rho_2, \mu \in \mathbb{R}$ and $q_1, q_2 \geq 2$ and let the following condition be satisfied*

$$\rho_1 - 1/q_1 = \mu = 1 - (\rho_2 - 1/q_2). \quad (6)$$

Then

$$1. \quad \|H(\cdot)U_0; L^{q_1}(\mathbb{R}, Y^{\rho_1})\| \leq C\|U_0; Y^\mu\|.$$

2. *For any interval $I = [0, T)$, $T \leq \infty$, then*

$$\|(\omega(\cdot), \omega_t(\cdot)); L^{q_1}(I, Y^{\rho_1})\| \leq C\|f; L^{\bar{q}_2}(I, H^{-\rho_2})\|.$$

The constants C are independent of the interval I .

For the proof of Theorem 2 we need the following proposition

Proposition 4 ([9, Lemma 3.4] and [13])

Let $1 \leq N \leq 3$, $f \in W^{1,\infty}(\mathbb{R}^N)$ and $g \in L^\infty(\mathbb{R}^N)$. Then, we have

$$\|\dot{K}(t)f; L^\infty(\mathbb{R}^N)\| \leq \max(1, t)\|f; W^{1,\infty}(\mathbb{R}^N)\|, \quad t > 0,$$

$$\|K(t)g; L^\infty(\mathbb{R}^N)\| \leq t\|g; L^\infty(\mathbb{R}^N)\|, \quad t > 0.$$

We will need also to deal with the Sobolev norm of a power of a function. The following is a direct consequence of [2]:

Proposition 5 Let $p \in \mathbb{N} \setminus \{0, 1\}$, $s > -N/2$, $s \neq N/2$ and $\nu(s, p) := \sup\{0, (N/2 - s)(p - 1)/p\}$. Then, for a nonnegative function $f \in H^{s+\nu(s, p)}(\mathbb{R}^N)$, $f^p \in H^s(\mathbb{R}^N)$ and there exists a positive constant C such that

$$\|f^p; H^s(\mathbb{R}^N)\| \leq C\|f; H^{s+\nu(s, p)}(\mathbb{R}^N)\|^p.$$

Next, we give the Gagliardo-Nirenberg inequality which is a direct consequence of [3, Theorem 1.3.4] (see also [4, Theorem 9.3]).

Proposition 6 (Gagliardo-Nirenberg)

Let q, r be such that $1 \leq q, r \leq \infty$, and let j, m be integers, $0 \leq j < m$. Let $a \in [j/m, 1]$ ($a < 1$ if $m - j - N/r$ is an integer ≥ 0), and let p be given by

$$\frac{1}{p} = \frac{j}{N} + a\left(\frac{1}{r} - \frac{m}{N}\right) + (1 - a)\frac{1}{q}.$$

For $f \in L^q(\mathbb{R}^N)$ such that $D^\alpha f \in L^r(\mathbb{R}^N)$ with $|\alpha| = m$, we have $D^\alpha f \in L^p(\mathbb{R}^N)$ with $|\alpha| = j$ and there exists a positive constant C such that

$$\sum_{|\alpha|=j} \|D^\alpha f\|_{L^p} \leq C \left(\sum_{|\alpha|=m} \|D^\alpha f\|_{L^r} \right)^a \|f\|_{L^q}^{1-a}.$$

For the second theorem we need

Proposition 7 Assume that $\mu \in (1, 2) \cup \mathbb{N}^*$ and $p \in \mathbb{R} \cap (1, \infty) \cap [\mu - 1, \infty)$. For a nonnegative function $f \in L^\infty(\mathbb{R}^N) \cap H^{\mu-1}(\mathbb{R}^N)$, $f^p \in H^{\mu-1}(\mathbb{R}^N)$ and there exists a positive constant C such that

$$\|f^p\|_{H^{\mu-1}(\mathbb{R}^N)} \leq C\|f\|_{L^\infty(\mathbb{R}^N)}^{p-1}\|f\|_{H^{\mu-1}(\mathbb{R}^N)}.$$

Proof. First case: $\mu \in (1, 2)$. Using the mean value theorem and the following equivalent norm of $\|\cdot\|_{H^{\mu-1}}$ (see [1, Theorem 7.48 page 214]), we

have:

$$\begin{aligned}
\|f^p\|_{H^{\mu-1}}^2 &:= \|f^p\|_{L^2}^2 + \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(f^p(x) - f^p(y))^2}{|x - y|^{N+2\mu-2}} dx dy \\
&= \|f^p\|_{L^2}^2 + \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(pz^{p-1}(f(x) - f(y)))^2}{|x - y|^{N+2\mu-2}} dx dy \\
&\leq p^2 \|f^{p-1}\|_{L^\infty}^2 \left[\|f\|_{L^2}^2 + \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(f(x) - f(y))^2}{|x - y|^{N+2\mu-2}} dx dy \right] \\
&\leq p^2 \|f\|_{L^\infty}^{2(p-1)} \|f\|_{H^{\mu-1}}^2,
\end{aligned}$$

where $\min(f(x), f(y)) < z < \max(f(x), f(y))$ for every $x, y \in \mathbb{R}^N$.

Second case: $\mu \in \mathbb{N}^*$. We have

$$\|f^p\|_{H^{\mu-1}} := \left(\sum_{0 \leq |\alpha| \leq \mu-1} \|D^\alpha f^p\|_{L^2}^2 \right)^{1/2}. \quad (7)$$

As

$$D^\alpha(f^p) = \sum_{|\beta_1| + \dots + |\beta_\alpha| = |\alpha|} C_{|\alpha|, |\beta_1|, \dots, |\beta_\alpha|} f^{p-|\alpha|} D^{\beta_1} f \dots D^{\beta_\alpha} f,$$

we conclude that

$$\begin{aligned}
\|D^\alpha f^p\|_{L^2} &\leq C \|f\|_{L^\infty}^{p-|\alpha|} \sum_{|\beta_1| + \dots + |\beta_\alpha| = |\alpha|} \|D^{\beta_1} f \dots D^{\beta_\alpha} f\|_{L^2} \\
&\leq C \|f\|_{L^\infty}^{p-|\alpha|} \sum_{|\beta_1| + \dots + |\beta_\alpha| = |\alpha|} \|D^{\beta_1} f\|_{L^{\frac{2|\alpha|}{|\beta_1|}}} \dots \|D^{\beta_\alpha} f\|_{L^{\frac{2|\alpha|}{|\beta_\alpha|}}}
\end{aligned}$$

thanks to Hölder's inequality. Using Proposition 6, we obtain

$$\begin{aligned}
\|D^\alpha f^p\|_{L^2} &\leq C \|f\|_{L^\infty}^{p-|\alpha|} \sum_{|\beta_1| + \dots + |\beta_\alpha| = |\alpha|} \left[\sum_{|\beta| = |\alpha|} \|D^\beta f\|_{L^2} \right]^{\frac{|\beta_1|}{|\alpha|}} \|f\|_{L^\infty}^{1-|\beta_1|/|\alpha|} \dots \\
&\quad \left[\sum_{|\beta| = |\alpha|} \|D^\beta f\|_{L^2} \right]^{\frac{|\beta_\alpha|}{|\alpha|}} \|f\|_{L^\infty}^{1-|\beta_\alpha|/|\alpha|} \\
&\leq C \|f\|_{L^\infty}^{p-1} \|D^\alpha f\|_{L^2},
\end{aligned}$$

which gives, using (7), the desired estimates. $\square$

3 Proof of theorems 1 and 2

Proof of Theorem 1. Without loss of generality, we may assume that the initial data (u_0, v_0) is of compact support. By finite speed of propagation, the solution (u, u_t) is also of compact support. This allows us to use

Proposition 3 on bounded intervals $[0, T]$ with usual Sobolev spaces instead of homogeneous Sobolev spaces (since for compactly supported distributions the norms are equivalent, see [16]).

Now, we write (1) as

$$\begin{cases} U_t - AU = F(U) \\ U(0, x) = U_0(x) \in Y^\mu, \end{cases} \quad (8)$$

where $U = (u, v)$, $A = \begin{bmatrix} 0 & I \\ \Delta & 0 \end{bmatrix}$ and $F(U) := \begin{bmatrix} 0 \\ v|v|^{p-1} \end{bmatrix}$. With theses notations, local existence for (1) is equivalent to local existence for (8), and this is equivalent to the following integral equation

$$\begin{cases} \text{Find } T > 0 \text{ and a unique solution,} \\ U \in C^0([0, T]; Y^\mu) \text{ of } U(t) = H(t)U_0 + L(U)(t), \end{cases} \quad (9)$$

where $H(t)U_0 = \begin{bmatrix} \dot{K}(t)u_0 + K(t)v_0 \\ \Delta K(t)u_0 + \dot{K}(t)v_0 \end{bmatrix}$ and $L(U)(t) = \begin{bmatrix} [K \star v^p](t) \\ [\dot{K} \star v^p](t) \end{bmatrix}$, where v^p denotes $v|v|^{p-1}$.

In order to use Fixed Point Theorem, let us introduce the following metric space

$$X := \{\phi = (\phi_1, \phi_2) \in L^q(0, T; Y^\rho) \text{ s.t. } \|\phi - HU_0; L^q(0, T; Y^\rho)\| \leq \lambda\}$$

where T and λ are positive constants and ρ and q satisfy (6). These constants will be fixed later.

For $\phi, \psi \in X$ denote by

$$\|\phi; X\| := \|\phi - HU_0; L^q(0, T; Y^\rho)\|$$

(HU_0 denotes the function $t \mapsto H(t)U_0$) and the natural induced distance

$$d(\phi, \psi) := \|\phi - \psi; L^q(0, T; Y^\rho)\|.$$

Finally define the map Φ on X by $\Phi(U)(t) := H(t)U_0 + L(U)(t)$.

First step: X is invariant under Φ . Let $U \in X$, by proposition 3 we have

$$\|\Phi(U); X\| \leq \|L(U); L^q(0, T; Y^\rho)\| \leq C\|v^p; L^{\bar{q}_2}(0, T; H^{-\rho_2})\|$$

for all (q_2, ρ_2) satisfying (6). Take $q_2 := +\infty$, hence $\rho_2 := 1 - \mu$. Then

$$\|\Phi(U); X\| \leq C\|v^p; L^1(0, T; H^{\mu-1})\|.$$

By proposition 5

$$\|v^p; H^{\mu-1}\| \leq C\|v; H^{\mu-1+\nu(\mu-1, p)}\|^p,$$

thus

$$\|\Phi(U); X\| \leq C\|v; L^p(0, T; H^{\mu-1+\nu(\mu-1, p)})\|^p. \quad (10)$$

If $\mu \geq 1 + N/2$ then $\nu(\mu - 1, p) = 0$ and choosing $\varepsilon = 1$, $q = \infty$ and $\rho = \mu$ then (10) gives

$$\begin{aligned} \|\Phi(U); X\| &\leq C\|v; L^p(0, T; H^{\mu-1})\|^p = C \int_0^T \|v; H^{\mu-1}\|^p \\ &\leq CT\|v; L^\infty(0, T; H^{\mu-1})\|^p \leq CT^\varepsilon \|U; L^q(0, T; Y^\rho)\|^p. \end{aligned} \quad (11)$$

If $\mu < 1 + N/2$, define $\varepsilon := 1 - p\nu(\mu - 1, p) = 1 - (p-1)(1 + N/2 - \mu) \in (0, 1)$ by (2). Then $\mu + \nu(\mu - 1, p) - \frac{1-\varepsilon}{p} = \mu$. Using Hölder's inequality we have

$$\|v; L^p(0, T; H^{\mu-1+\nu(\mu-1, p)})\|^p \leq CT^\varepsilon \|v; L^{\frac{p}{1-\varepsilon}}(0, T; H^{\mu-1+\nu(\mu-1, p)})\|^p.$$

By choosing $\rho = \mu + \nu(\mu - 1, p)$ and $q = p/(1 - \varepsilon)$, and using proposition 3, inequality (10) gives

$$\|\Phi(U); X\| \leq CT^\varepsilon \|v; L^q(0, T; H^{\rho-1})\| \leq CT^\varepsilon \|U; L^q(0, T; Y^\rho)\|^p.$$

We see that, in both cases one has

$$\|\Phi(U); X\| \leq CT^\varepsilon \|U; L^q(0, T; Y^\rho)\|^p, \quad (12)$$

with ρ and q satisfying (6). Thus we have

$$\|\Phi(U); X\| \leq CT^\varepsilon \|U; L^q(0, T; Y^\rho)\|^p \leq CT^\varepsilon [\|U; X\| + \|HU_0; L^q(0, T; Y^\rho)\|]^p,$$

and, as ρ and q satisfy (6), by proposition 3 we have

$$\|\Phi(U); X\| \leq CT^\varepsilon [\|U; X\| + \|U_0; Y^\mu\|]^p.$$

Therefore, X is invariant by Φ if T and λ are such that

$$CT^\varepsilon (\lambda + \|U_0; Y^\mu\|)^p \leq \lambda. \quad (13)$$

Second step: Φ is a contraction on X .

This is mainly the same ideas. Let $U, V \in X$. Then

$$d(\Phi(U), \Phi(V)) = \|L(U) - L(V); L^q(0, T; Y^\rho)\| \leq C\|U_2^p - V_2^p; L^{\overline{q_2}}(0, T; H^{-\rho_2})\|$$

for ρ_2 and q_2 satisfying (6). Take $q_2 := +\infty$, we get $\rho_2 := 1 - \mu$ and then

$$d(\Phi(U), \Phi(V)) \leq C\|U_2^p - V_2^p; L^1(0, T; H^{\mu-1})\|.$$

Now, we write $U_2^p - V_2^p = (U_2 - V_2)P(U_2, V_2)$ where P is a homogeneous polynomial of degree $p - 1$. Using proposition 5 and the convexity of the exponential function to obtain

$$\begin{aligned} d(\Phi(U), \Phi(V)) &\leq C\|U_2 - V_2; L^p(0, T; H^{\mu-1+\nu(\mu-1, p)})\| \\ &\quad \times \left[\|U_2; L^p(0, T; H^{\mu-1+\nu(\mu-1, p)})\|^{p-1} + \|V_2; L^p(0, T; H^{\mu-1+\nu(\mu-1, p)})\|^{p-1} \right]. \end{aligned}$$

By the same analysis, and a similar calculation as in the first step, we get

$$d(\Phi(U), \Phi(V)) \leq Cd(U, V)T^\varepsilon[\lambda + \|U_0; Y^\mu\|]^{p-1}.$$

Finally, Φ is a contraction on X if λ and T satisfy

$$CT^\varepsilon[\lambda + \|U_0; Y^\mu\|]^{p-1} < 1. \quad (14)$$

By choosing λ and T satisfying (13) and (14), Fixed Point Theorem ensures existence and uniqueness.

Third step: continuity of the solution.

We have obtained existence and uniqueness of a solution $U = (u, u_t) \in X \subset L^q(0, T; Y^\rho)$ where ρ and q satisfy (6). Let's show that $U \in L^{q'}(0, T; Y^{\rho'})$ for any ρ' and q' satisfying (6). This is a similar calculation to the first step. Indeed,

$$\begin{aligned} \|U; L^{q'}(0, T; Y^{\rho'})\| &\leq C\|v^p; L^1(0, T; Y^{\mu-1})\| + \|U_0; Y^\mu\| \\ &\leq CT^\varepsilon\|U; L^q(0, T; Y^\rho)\|^p + \|U_0; Y^\mu\| < \infty. \end{aligned}$$

In particular $U \in L^\infty(0, T; Y^\mu)$. Now, using (9), for $0 \leq t_0 \leq t$, we have

$$\|U(t) - U(t_0); L^\infty(t_0, t; Y^\mu)\| \leq \|U(t) - H(t - t_0)U_0\| + \|U(t_0) - H(t - t_0)U_0\|.$$

As in the first step

$$\|U(t) - H(t - t_0)U_0; L^\infty(t_0, t; Y^\mu)\| \leq C(t - t_0)^\varepsilon\|U(t); L^q(t_0, t; Y^\rho)\|^p,$$

and the strong continuity of the C_0 -group H associated to the wave equation implies that

$$\lim_{t \rightarrow t_0} \|H(t - t_0)U_0 - U_0; L^\infty(t_0, t; Y^\mu)\| = 0.$$

Therefore, $U \in C^0([0, T]; Y^\mu)$. □

Proof of Theorem 2. Using Proposition 4, the proof is similar to that of Theorem 1 where we use Proposition 7 instead of Proposition 5. □

References

- [1] Robert A. Adams. *Sobolev spaces*. Academic Press [A subsidiary of Harcourt Brace Jovanovich, Publishers], New York-London, 1975. Pure and Applied Mathematics, Vol. 65.
- [2] A. Bachelot. Formes quadratiques compatibles dans les espaces de type l^p . *Compte-rendu CNRS*, (8406), 1984.

- [3] Thierry Cazenave and Alain Haraux. *Introduction aux problèmes d'évolution semi-linéaires*, volume 1 of *Mathématiques & Applications (Paris) [Mathematics and Applications]*. Ellipses, Paris, 1990.
- [4] Avner Friedman. *Partial differential equations of parabolic type*. Prentice-Hall Inc., Englewood Cliffs, N.J., 1964.
- [5] Vladimir Georgiev and Grozdena Todorova. Existence of a solution of the wave equation with nonlinear damping and source terms. *J. Differential Equations*, 109(2):295–308, 1994.
- [6] J. Ginibre and G. Velo. Generalized Strichartz inequalities for the wave equation. *J. Funct. Anal.*, 133(1):50–68, 1995.
- [7] A. Haraux. Remarks on the wave equation with a nonlinear term with respect to the velocity. *Portugal. Math.*, 49(4):447–454, 1992.
- [8] A. Haraux and E. Zuazua. Decay estimates for some semilinear damped hyperbolic problems. *Arch. Rational Mech. Anal.*, 100(2):191–206, 1988.
- [9] Takafumi Hosono and Takayoshi Ogawa. Large time behavior and L^p - L^q estimate of solutions of 2-dimensional nonlinear damped wave equations. *J. Differential Equations*, 203(1):82–118, 2004.
- [10] M. Kopáčeková. Remarks on bounded solutions of a semilinear dissipative hyperbolic equation. *Comment. Math. Univ. Carolin.*, 30(4):713–719, 1989.
- [11] Frank Merle and Hatem Zaag. Determination of the blow-up rate for the semilinear wave equation. *Amer. J. Math.*, 125(5):1147–1164, 2003.
- [12] Salim A. Messaoudi. Blow up and global existence in a nonlinear viscoelastic wave equation. *Math. Nachr.*, 260:58–66, 2003.
- [13] Kenji Nishihara. L^p - L^q estimates of solutions to the damped wave equation in 3-dimensional space and their application. *Math. Z.*, 244(3):631–649, 2003.
- [14] Jalal Shatah and Michael Struwe. *Geometric wave equations*. 2:viii+153, 1998.
- [15] Robert S. Strichartz. Restrictions of Fourier transforms to quadratic surfaces and decay of solutions of wave equations. *Duke Math. J.*, 44(3):705–714, 1977.
- [16] Hans Triebel. *Theory of function spaces*, volume 78 of *Monographs in Mathematics*. Birkhäuser Verlag, Basel, 1983.